

The disruption time of clusters in selected regions of four galaxies

Henny J.G.L.M. Lamers^{1,2} and Stratos G. Boutloukos^{1,3,4}

¹ Astronomical Institute, Utrecht University, Princetonplein 5, NL-3584CA, Utrecht, The Netherlands

² SRON Laboratory for Space Research

³ Aristotle University, Thessaloniki, Greece

⁴ University of Tuebingen, Institute of Theoretical Astrophysics and Computational Physics, Auf der Morgenstelle 10, D-72076, BRD

Abstract. If the cluster formation rate is constant and the disruption time of clusters depends on their initial mass as $t_{\text{dis}} = t_4 \times (M_{\text{cl}}/10^4 M_{\odot})^{\gamma}$, the values of t_4 and γ can be derived in a very simple way from the age and mass histograms of large homogeneous samples of clusters with reliable age and/or mass determinations. We demonstrate the method and derive the values of t_4 and γ from cluster samples in selected fields of four galaxies: M51, M33, SMC and the solar neighbourhood. The values of γ are the same within their uncertainties in the four galaxies with $\langle \gamma \rangle = 0.62 \pm 0.06$. However, the disruption time t_4 of clusters of $10^4 M_{\odot}$ is very different in the different locations. It is shortest in the inner region of M51 ($t_4 = 4 \times 10^7$ yrs) and highest in the SMC ($t_4 = 4 \times 10^9$ yrs).

1 Introduction

Studies of the cluster formation history in different galaxies, based on *observed* cluster samples, rarely take into account the fact that clusters may have been disrupted. We have started a series of studies to disentangle the effects of cluster formation and cluster disruption. Here we present the method for the determination of the disruption time in galaxies with a nearly constant cluster formation rate, with an application to selected regions in four galaxies.

Oort [12] noticed the lack of old open clusters in the solar neighbourhood and derived statistically that Galactic clusters disrupt on a timescale of 5×10^8 years. Several authors have found that the mean age of clusters in the LMC and SMC is larger than that of clusters in the solar neighbourhood and concluded that the disruption time must be longer in the MCs (e.g. [9]). Actually, the disruption time of clusters is expected to depend on their mass and on the local conditions in their host galaxy. Modern observations of large homogeneous cluster samples, e.g. with *HST*, enable the determination of the cluster disruption time in different galaxies.

The disruption time of clusters is expected to scale with the relaxation time

$$t_{\text{dis}} \sim t_{\text{rxt}} \sim R_{\text{cl}}^{3/2} M_{\text{cl}}^{1/2} \quad (1)$$

where R_{cl} is the cluster radius and M_{cl} is the cluster mass (see e.g. [13],[14]). If all clusters in a limited region of a galaxy had the same radius, then Eq.

(1) would imply that $t_{\text{dis}} \sim M_{\text{cl}}^{1/2}$. If the cluster radius depends on its mass as $R_{\text{cl}} = C_{\text{gal}} \times M_{\text{cl}}^\delta$, then we expect $t_{\text{dis}} \sim C_{\text{gal}}^{3/2} M_{\text{cl}}^\gamma$ with $\gamma = (1 + 3\delta)/2$, where C_{gal} is a constant that depends on the ambient pressure in the region of the galaxy where the clusters are located. (Obviously, this expected relation does not hold for clusters in highly elliptical orbits.)

In this paper we derive the empirical relation between the disruption time and the mass of clusters in different locations of galaxies, based on the mass and/or age distributions of magnitude limited cluster samples.

2 The predicted mass and age distributions of surviving clusters above a given magnitude limit.

Suppose that:

1. the *disruption time of clusters* depends on their initial mass M_{cl} as

$$t_{\text{dis}} = t_4 \times (M_{\text{cl}}/10^4 M_\odot)^\gamma \quad (2)$$

where t_4 is the disruption time of a cluster with an initial mass of $10^4 M_\odot$,

2. the *formation rate of clusters* is constant,
3. the *initial mass function of clusters* can be written as $N(M_{\text{cl}}) \sim M_{\text{cl}}^{-\alpha}$, with $\alpha \simeq 2$ [16],
4. *clusters fade below the detection limit*, in the wavelength band that determines the magnitude limit of the cluster sample, due to the evolution of their stars as $F_\lambda \sim t^{-\zeta}$. The value of ζ can be derived from cluster evolution models, e.g. the Starburst99 models of Leitherer et al. [11].

Then it is easy to show that both the age and mass distributions of surviving clusters above a certain magnitude limit will consist of double power laws (shown in Fig. 1) of the type:

- $dN_{\text{cl}}/dM_{\text{cl}} \sim M_{\text{cl}}^{(1/\gamma)-\alpha}$ for low mass clusters due to fading
- $dN_{\text{cl}}/dM_{\text{cl}} \sim M_{\text{cl}}^{\gamma-\alpha}$ for high mass clusters due to disruption
- $dN_{\text{cl}}/dt \sim M_{\text{cl}}^{\zeta(1-\alpha)}$ for young clusters due to fading
- $dN_{\text{cl}}/dt \sim M_{\text{cl}}^{(1-\alpha)/\gamma}$ for old clusters due to disruption

These relations are derived by Boutloukos & Lamers [4] (hereafter BL02).

The vertical shift of the fading lines in Fig. 1 depends on the detection limit of the observations and on the number of clusters formed per unit time in the region of the host galaxy that is studied. The horizontal shift of the disruption line depends on t_4 . The slope of the disruption line depends on the values of γ and α . So both t_4 and γ can be derived from the cluster statistics in a simple straightforward way.

These double power laws are strictly valid if the clusters disrupt instantaneously when they reach an age of t_{dis} . However, it can be shown that gradual

cluster disruption results in similar power law distributions (BL02). In that case the transition between the two powerlaws is not a sharp bend, but it is more gradual. This is demonstrated in Fig. 2. that shows the predicted mass and age distributions in case the cluster mass decreases gradually on a timescale that depends on the mass as in Eq. (2). The full lines are the predicted relations for gradual disruption, whereas the dashed lines are the relations in case of instantaneous disruption. They show a similar double power law as in Fig. 1, but with a gradual transition.




Fig. 1. Schematic pictures of the predicted mass distribution (left) and age distribution (right) of clusters due to fading (full lines) and disruption (dotted lines). The y -axes have arbitrary units. (figure from BL02)




Fig. 2. Predictions for gradual decay of clusters with an age dependent detection limit and with a disruption time that depends on the actual cluster mass as given by Eq. (2). Left: the predicted mass distribution (full line). Right: the predicted age distribution (full line). The dash-dotted lines show the prediction if there was no disruption but only evolutionary fading. The dotted lines show the prediction for evolutionary fading and instantaneous disruption. The parameters differ from those used in Fig. 1. (figure from BL02)

3 Application to regions in four galaxies

We have applied this method to samples of clusters observed with *HST* in the inner region of M51 by Bik et al. [3], in M33 by Chandar et al. [5],[6],[7] and to the cluster samples of the SMC by Hodge [10] and of the solar neighbourhood by Wielen [15]. The distributions indeed show the power-law decrease at high mass and/or high age due to disruption. Figures 3a shows the mass distribution of the M51 cluster sample and Fig. 3b shows the age distribution of the M33 cluster sample. (BL02 also studied the age distribution of clusters in M51 and the mass distribution of clusters in M33.) The full lines are the predicted distributions for evolutionary fading below the detection limit for these samples. (The wiggles at low age or small mass are due to the appearance and disappearance of red supergiants.) At high age or high mass the observed distributions are powerlaws that clearly decrease much steeper than predicted for evolutionary fading only. This is due to disruption. Figures 4a and 4b show the age distributions of clusters in the SMC and the solar neighbourhood respectively. The age distribution of the clusters in the SMC and the solar neighbourhood show a flat part at young ages, because the clusters do not fade below the detection limit in the samples used. Therefore, the distributions at young age show the constant cluster formation rate. From the observed powerlaw relations we derived the values of the disruption parameters t_4 and γ .

The results are summarized in Table 1. The values of γ are very similar (within the uncertainties) for the cluster samples in the four galaxies, with a mean value of $\langle \gamma \rangle = 0.62 \pm 0.06$. The values of t_4 , however, are very different for the different galaxies: the inner region of M51 has the shortest cluster disruption time and the SMC has the longest disruption times. The difference amounts to about a factor 10^2 .

Table 1. The parameters of the disruption time: $t_{\text{dis}} = t_4 \times (M_{\text{cl}}/10^4)^\gamma$

Galaxy	Region	Nr	Age range	Mass range	$\log t_4$	γ
	$r_{\text{gal}}(\text{kpc})$	clusters	$\log(\text{yrs})$	$\log M_\odot$	$\log(\text{yrs})$	
M51	0.8 - 3.1	512	6.0 - 9.7	3.0 - 5.2	7.64 ± 0.22	0.57 ± 0.10
M33	0.8 - 5.0	49	6.5 - 10.0	3.6 - 5.6	8.12 ± 0.30	0.72 ± 0.12
MW	7.5 - 9.5	72	7.2 - 10.0	—	9.0 ± 0.3	0.60 ± 0.12
SMC	0 - 2	314	7.6 - 10.0	—	9.6 ± 0.3	0.61 ± 0.08
Mean						0.62 ± 0.06




Fig. 3. The mass distribution of M51 clusters in the inner 0.8 - 3.1 kpc (left) and the age distribution of M33 clusters at $0.8 < r_{\text{Gal}} < 5.0$ kpc (right). The full lines are the predicted decrease due to fading below the detection limit (from Starburst99 models). The dashed lines are powerlaw fits for disruption. (figure from BL02)




Fig. 4. The age distributions of clusters in the SMC (left) and in the solar neighbourhood (right). Evolutionary fading below the detection limit is not important for these cluster samples, so the young clusters show a constant cluster formation rate. The dashed lines are powerlaw fits for disruption. (figure from BL02)

4 Future studies

The full study has been described by BL02. Future planned or ongoing studies include:

- study of the disruption time of clusters in different regions of M51 (Bastian & Lamers, [1])
- study of the disruption time of clusters in the starburst galaxy M82 (de Grijs et al. [8])
- study of the disruption time of clusters in M81 as a function of mass and radius (Bastian et al., [2])
- comparison of the derived disruption times with N-body simulations (Lamers & Portegies Zwart, in preparation)
- comparison of the disruption times of interacting and non-interacting galaxies (Bastian et al., in preparation)

Acknowledgements

We thank Arjan Bik and Nate Bastian for their contribution to the analysis of the M51 *HST*-data. The stay of Stratos Boutloukos at the Astronomical Institute in Utrecht was supported by a grant from the Erasmus student project of the European Community.

References

1. N. Bastian, H.J.G.L.M. Lamers: these proceedings (2003)
2. N. Bastian, R. van Deelen, H.J.G.L.M. Lamers, R. de Grijs, O. Noor, M. Bosman: (in preparation) (2003)
3. A. Bik, H.J.G.L.M. Lamers, N. Bastian, N. Panagia, M. Romaniello: A&A (in press) (2002)
4. S.G. Boutloukos, H.J.G.L.M. Lamers: MNRAS (in press) (2002)
5. R. Chandar, L. Bianchi, H.C. Ford: ApJ, **517**, 668 (1999a)
6. R. Chandar, L. Bianchi, H.C. Ford: ApJS **122**, 431 (1999b)
7. R. Chandar, L. Bianchi, H.C. Ford, B. Salasnich: PASP, **111**, 794 (1999c)
8. R. de Grijs, N. Bastian, H.J.G.L.M. Lamers: MNRAS (submitted) (2002)
9. R.A.W. Elson, S.M. Fall: ApJ, **299**, 211 (1985)
10. P. Hodge: PASP, **99**, 742 (1987)
11. C. Leitherer, D. Schaerer, J.D. Goldader, et al.: ApJS, **123**, 3 (1999)
12. J.H. Oort: *Stellar Populations*, Ric. Astr. Specolo Astr. Vaticana, **5**, 63 (1958)
13. L. Spitzer Jr: ApJ **127**, 544 (1958)
14. L. Spitzer Jr: *Dynamical Evolution of Globular Clusters*, (Princeton Univ. Press, Princeton 1987)
15. R. Wielen: A&A, **13**, 309 (1971)
16. Q. Zhang, S.M. Fall: ApJ, **527**, L81 (1999)